Disease Case Classification PHAST Measures of Local Public Health Services and Activities 8/07/2017

Disease	Suspected case classification	Probable case classification	Confirmed case classification	Source of classification (URL of CDC page)
Rubella	Any generalized rash illness of acute onset that does not meet the criteria for probable or confirmed rubella or any other illness.	In the absence of a more likely diagnosis, an illness characterized by all of the following: acute onset of generalized maculopapular rash; and temperature greater than 99.0° F or 37.2° C, if measured; and arthralgia, arthritis, lymphadenopathy, or conjunctivitis; and lack of epidemiologic linkage to a laboratory-confirmed case of rubella; and noncontributory or no serologic or virologic testing.	A case with or without symptoms who has laboratory evidence of rubella infection confirmed by one or more of the following laboratory tests: isolation of rubella virus; or detection of rubella-virus specific nucleic acid by polymerase chain reaction; or IgG seroconversion or a significant rise between acute- and convalescent-phase titers in serum rubella IgG antibody level by any standard serologic assay (not explained by MMR vaccination during the previous 6-45 days); or positive serologic test for rubella IgM antibody (not explained by MMR vaccination during the previous 6-45 days and not otherwise ruled out by more specific testing in a public health laboratory) or an illness characterized by all of the following: acute onset of generalized maculopapular rash; and temperature greater than 99.0°F or 37.2°C; and arthralgia, arthritis, lymphadenopathy, or conjunctivitis; and epidemiologic linkage to a laboratory-confirmed case of rubella.	http://wwwn.cdc.gov/NNDSS/script/casedef .aspx?CondYrID=909&DatePub=1/1/2013% 2012:00:00%20AM
Measles		In the absence of a more likely diagnosis, an illness that meets the clinical description (An acute illness characterized by: generalized, maculopapular rash lasting ≥3 days; and temperature ≥101°F or 38.3°C; and cough, coryza, or conjunctivitis) with: no epidemiologic linkage to a laboratory-confirmed measles case; and noncontributory or no measles laboratory testing.	An acute febrile rash illness (temperature does not need to reach ≥101°F/38.3°C and rash does not need to last ≥3 days) with: isolation of measles virus from a clinical specimen (not explained by MMR vaccination during the previous 6-45 days); or IgG seroconversion (not explained by MMR vaccination during the previous 6-45 days) or a significant rise in measles immunoglobulin G antibody (not explained by MMR vaccination during the previous 6-45 days) using any evaluated and validated method; or a positive serologic test for measles immunoglobulin M antibody (not explained by MMR vaccination during the previous 6-45 days; not otherwise ruled out by other confirmatory testing or more specific measles testing in a public health laboratory); or direct epidemiologic linkage to a case confirmed by one of the methods above.	http://wwwn.cdc.gov/NNDSS/script/casedef .aspx?CondYrID=908&DatePub=1/1/2013% 2012:00:00%20AM
Congenital rubella	An infant that does not meet the criteria for a probable or confirmed case but who has one of more of the following clinical findings: cataracts or congenital glaucoma, congenital heart disease (most commonly patent ductus arteriosus or peripheral pulmonary artery stenosis), hearing impairment, pigmentary retinopathy, purpura, hepatosplenomegaly, jaundice, microcephaly, developmental delay, meningoencephalitis, OR radiolucent bone disease.	An infant without an alternative etiology that does not have laboratory confirmation of rubella infection but has at least 2 of the following*: cataracts or congenital glaucoma,*congenital heart disease (most commonly patent ductus arteriosus or peripheral pulmonary artery stenosis), hearing impairment, OR pigmentary retinopathy; OR An infant without an alternative etiology that does not have laboratory confirmation of rubella infection but has at least one or more of the following: cataracts or congenital glaucoma,*congenital heart disease (most commonly patent ductus arteriosus or peripheral pulmonary artery stenosis), hearing impairment, OR pigmentary retinopathy AND one or more of the following: purpura, hepatosplenomegaly, jaundice, microcephaly, developmental delay, meningoencephalitis, OR radiolucent bone disease. *In probable cases, either or both of the eye-related findings (cataracts and congenital glaucoma) count as a single complication. In cases classified as infection only, if any compatible signs or symptoms (e.g., hearing loss) are identified later, the case is reclassified as confirmed.		_aspx?CondYrID=834&DatePub=1/1/2010% 2012:00:00%20AM
Mumps	Parotitis, acute salivary gland swelling, orchitis, or oophoritis unexplained by another more likely diagnosis, or a positive lab result with no mumps clinical symptoms (with or without epidemiological-linkage to a confirmed or probable case).	Acute parotitis or other salivary gland swelling lasting at least 2 days, or orchitis or oophoritis unexplained by another more likely diagnosis, in: a person with a positive test for serum anti-mumps immunoglobulin M (IgM) antibody, or a person with epidemiologic linkage to another probable or confirmed case or linkage to a group/community defined by public health during an outbreak of mumps.	A positive mumps laboratory confirmation for mumps virus with reverse transcription polymerase chain reaction (RT-PCR) or culture in a patient with an acute illness characterized by any of the following: acute parotitis or other salivary gland swelling, lasting at least 2 days, aseptic meningitis, encephalitis, hearing loss, orchitis, oophoritis, mastitis, pancreatitis.	http://wwwn.cdc.gov/NNDSS/script/casedef _aspx?CondYrID=783&DatePub=1/1/2012% 2012:00:00%20AM

		to the absence of a serial likely discussion on the illness with according	There is no definition for "confirmed" tetanus.	hater // and a ser /NNIDCS /a-si-t/d-f
		In the absence of a more likely diagnosis, an acute illness with muscle spasms or hypertonia, and diagnosis of tetanus by a health care provider; or		http://wwwn.cdc.gov/NNDSS/script/casedef
Tetanus		, , , , , , , , , , , , , , , , , , , ,		
		death, with tetanus listed on the death certificate as the cause of death or a		2012:00:00%20AM
		significant condition contributing to death.		
	A case of postdiarrheal HUS or identification of Shiga	A case with isolation of <i>E. coli</i> O157 from a clinical specimen, without	A case that meets the confirmed laboratory criteria for diagnosis. When available,	http://wwwn.cdc.gov/NNDSS/script/casedef
	toxin in a specimen from a clinically compatible case	confirmation of H antigen or Shiga toxin production; or a clinically	O and H antigen serotype characterization should be reported.	.aspx?CondYrID=951&DatePub=1/1/2014%
E. coli, shiga toxin	without the isolation of STEC.	compatible case who is a contact of an STEC case or is a member of a		2012:00:00%20AM
producing strains only		defined risk group during an outbreak; or identification of an elevated		
		antibody titer to a known STEC serotype from a clinically compatible case.		
	A case that meets the suspect laboratory criteria for	A clinically compatible case that is epidemiologically linked to a confirmed	A case that meets the confirmed laboratory criteria for diagnosis. When available,	http://wwwn.cdc.gov/NNDSS/script/casedef
Salmonellosis	diagnosis.	case, i.e., a contact of a confirmed case or member of a risk group as	O and H antigen serotype characterization should be reported.	.aspx?CondYrID=844&DatePub=1/1/2012%
	ulagriosis.	defined by public health authorities during an outbreak.	and it antigen service characterization should be reported.	2012:00:00%20AM
	A case that meets the suspect laboratory criteria for	A clinically compatible case that is epidemiologically linked to a confirmed	A case that meets the confirmed laboratory criteria for diagnosis.	http://wwwn.cdc.gov/NNDSS/script/casedef
Campylobacteriosis	diagnosis.	case of campylobacteriosis.	, ,	.aspx?CondYrID=627&DatePub=1/1/2012%
				2012:00:00%20AM
	A case that meets the suspect laboratory criteria for	A clinically compatible case that is epidemiologically linked, i.e., is a contact	A case that meets the confirmed laboratory criteria for diagnosis. When available,	http://wwwn.cdc.gov/NNDSS/script/casedef
Shigellosis	diagnosis.	of a confirmed case or a member of a risk group defined by public health	O antigen serotype characterization should be reported.	.aspx?CondYrID=849&DatePub=1/1/2012%
		authorities during an outbreak.		2012:00:00%20AM
		A clinically compatible case who had a consistent exposure (consumption of	A clinically compatible case with toxin detected in an epidemiologically implicated	http://epi.publichealth.nc.gov/cd/lhds/man
Ciguatera		fish such as barracuda, grouper, amberjack, and snapper).	fish.	uals/cd/casedefs/FOODBORNE POISONING
				CIGUATERA CD.pdf
Paralytic shellfish		A clinically compatible case that is not laboratory confirmed and not	A case that is laboratory confirmed, or that meets the clinical case definition, is	http://www.doh.wa.gov/Portals/1/Docume
· · · · · · · · · · · · · · · · · · ·		epidemiologically linked to a confirmed case.	not laboratory confirmed, and is epidemiologically linked to a confirmed case.	nts/5100/420-077-Guideline-
poisoning				ShellfishPoisoning.pdf
		A clinically compatible case with consumption of fish such as tuna,	A clinically compatible case with histamine detection in an epidemiologically	http://epi.publichealth.nc.gov/cd/lhds/man
Scombroid		mackerel, skipjack, bonito, mahi mahi, and blue fish within three hours of	implicated fish case that is epidemiologically linked to a confirmed case.	uals/cd/casedefs/FOODBORNE POISONING
		onset of symptom.		SCOMBROID CD.pdf
Mushroom poisoning			A clinically compatible case in someone with mushroom exposures	http://epi.publichealth.nc.gov/cd/lhds/man
				uals/cd/casedefs/FOODBORNE_POISONING
				_MUSHROOM_CD.pdf
		A clinically compatible case with an epidemiological link (e.g., ingestion of a	, ,	http://wwwn.cdc.gov/NNDSS/script/casedef
Botulism		home-canned food within the previous 48 hours)	persons who ate the same food as persons who have laboratory-confirmed	.aspx?CondYrID=622&DatePub=1/1/2011%
			botulism.	2012:00:00%20AM